BARC submission

Barau's Petrel – Bremer Canyon, Western Australia, 24th January, 2021.

Steve Reynolds & Carl Billingham

(BARC submission prepared on observer's behalf by Daniel Mantle & Plaxy Barratt

<u>Submission note:</u> although still a very rare bird elsewhere in Australia, Barau's Petrels have been recorded annually (January–March) off the southern coast of Western Australia since 2015. Tracking data also suggests they are likely regular off the NW of Western Australia.

<u>Circumstances of sighting:</u> a single Barau's Petrel *Pterodroma baraui* was spotted and photographed on the 24th January 2021 by Steve Reynolds during the Naturaliste Charter's Orca tour to the Bremer Canyon, southern coast of Western Australia.

<u>Description:</u> A long and slender-winged, medium-sized *Pterodroma* with mostly white underparts with distinctive black carpal bars, a longish graduated tail, and a distinctive darker cap. From the photo, the bird looks obviously longer winged than a Cookilaria petrel.

<u>Upperparts:</u> The upper wings are very dark grey to blackish and with little or no contrast with mantle or coverts evident in the images. The upper tail is only partly visible but appears dark grey/black and uniform in colour.

<u>Underparts</u>: The underwings are mostly white, framed by a well-defined black trailing edge, mostly black primaries and with very substantial black carpal bar that runs from the base of the primaries to the carpal joint and then diagonally back towards the body. The undertail coverts are white and the partly visible undertail feathers appear dark grey.

<u>Head</u>: A distinctive blackish cap extending below the eyes grades into smudgy pale grey neck sides. These smudgy grey neck sides extend into a ragged grey half-collar. The throat and chin are white, as are the lores and forehead and contrast markedly with the blackish cap.

Bare parts: moderately stout, all dark bill. Feet and legs not apparent in photos.

Distinguishing similar species:

White-necked Petrel *Pterodroma cervicalis*: any differences in jizz are not readily discernible from these photographs, although the bird does appear a little slim for a typical White-necked Petrel. The lack of a white hind neck is the main feature from these images to rule out White-necked Petrel. The minor differences in underwing pattern can't be easily distinguished from these photos. Vanuatu Petrel *Pterodroma occulta* is extremely similar but marginally smaller than White-necked Petrel and is distinguished from Barau's Petrel by the same plumage features.

Juan Fernandez Petrel *Pterodroma externa*: again, structurally quite similar without easily discernible differences from the photographic evidence presented herein. However, distinctive plumage differences include the much-reduced black carpal bar and thinner black trailing edge that would be expected for a Juan Fernandez Petrel.

All the smaller cookilaria petrels can be separated on structural jizz alone, even from these photographs. They are all substantially smaller with shorter, broader wings with more rounded hands. The thick black carpal bar also eliminates many of the cookilaria with reduced carpal ticks (e.g., Cook's and Stejneger's Petrel) whilst other regular Australian cookilaria such as Gould's and Black-winged Petrel may both be distinguished by differing head/neck patterns (Gould's is distinctively 'hooded', whilst Black-winged typically lacks the dark cap and often has a more pronounced grey half collar). The latter also has notably thicker black carpal bars and trailing edge to the underwings.

<u>Observer's experience</u>: Steve has seen multiple Barau's Petrels at the Bremer Canyon and at the Cocos islands and is very familiar with the typical seabirds of the region and any possible Australian confusion species. Carl is also experienced with many of the possible confusion species in Australia, though this was his first Barau's Petrel.

References:

CARTER, M., T. REID & P. LANSLEY (1989). Barau's Petrel *Pterodroma baraui*: A new species for Australia. Australian Bird Watcher 13: 39-43.

HOWELL, S.N.G. (2012). Petrels, Albatrosses, and Storm-Petrels of North America: A Photographic Guide. Princeton.

MARCHANT, S. & HIGGINS, P.J. (eds) (1990). Handbook of Australian, New Zealand and Antarctic Birds. Volume 1: Ratites to Ducks. Melbourne.

MENKHORST, P., ROGERS, D., CLARKE, R., DAVIES, J., MARSACK, P. & FRANKLIN, K. (2017): The Australian Bird Guide. CSIRO publishing.

ONLEY, D. & SCOFIELD, P. (2007): Albatrosses, Petrels & Shearwaters of the World. Princeton.

PINET, P., JAQUEMET, S., PINAUD, D., WEIMERSKIRCH, H., PHILLIPS, R.A., & LE CORRE, M. (2011). Migration, wintering distribution and habitat use of an endangered tropical seabird, Barau's Petrel *Pterodroma baraui*. Marine Ecology Progress Series 423: 291-302.



Images – Steve Reynolds, Bremer Canyon, WA, 24th January 2021